A photograph of a zebra finch perched on a dark brown tree branch. The bird has a grey body with a prominent yellow patch on its wing and a red beak. It is set against a dark background filled with numerous out-of-focus, glowing circular lights in various colors including green, red, blue, and yellow, creating a bokeh effect.

Dynamical systems and AI to model complex systems

Gabriel Mindlin

### The rationale for the course:

One of the crucial intellectual decisions facing young scientists today is determining the balance between **interpretable** and **data-driven** approaches when addressing a scientific problem.

## The rationale for the course:

One of the crucial intellectual decisions facing young scientists today is determining the balance between **interpretable** and **data-driven** approaches when addressing a scientific problem.

### **Interpretable science**

Variables have meaning

The relationship between the  
Variables are interpretable mechanisms

## The rationale for the course:

One of the crucial intellectual decisions facing young scientists today is determining the balance between **interpretable** and **data-driven** approaches when addressing a scientific problem.

### Interpretable science

Variables have meaning

The relationship between the  
Variables are interpretable mechanisms


### Data driven science

A complex "non interpretable" model  
(as a neural network) is trained with  
examples, and the model does not  
share with us the rationale behind its success

## Counterpoint with history



Kepler (1571-1630)



Newton (1643-1727)



Laplace (1749-1827): this is the “world’s system”  
All we have to do is to compute “f”, and the initial conditions

Why should we try to describe a problem with differential equations?  
Should we? Can we? **After all...**


Why should we try to describe a problem with differential equations?  
Should we? Can we? **After all...**

One of the biggest revolutions in science was data driven.  
And the “conversation” between Kepler and Newton is  
the first battle between data science and interpretable science

## Kepler's Laws


### First Law

All planets move around the sun in elliptical orbits with the sun at one of the foci




### Second Law

A planet sweeps out equal areas in equal intervals of time



### Third Law

The square of the orbital period of a planet is proportional to the cube of the orbit's semi-major axis



$$T^2 \propto a^3$$

T = Time to complete orbit  
a = Length of semi-major axis

The variables in Kepler's laws are interpretable, but we have no interpretable mechanism relating them.

## Newton's approach gave us more

We define the velocity.

$$\frac{dx}{dt} = v$$

$$\frac{dv}{dt} = \begin{cases} 0 & \text{In the absence of interactions with the universe (Galileo)} \\ \frac{1}{m} & \text{Multiplied by a functional form describing the interaction} \end{cases}$$

Some functions, fundamental ones

$$\mathbf{F} = \frac{-G m M}{r_{12}^2} \mathbf{e}_{12}$$

Others, phenomenological

$$F = -kx$$

## Newton's approach gave us more

We define the velocity.

$$\frac{dx}{dt} = v$$

$$\frac{dv}{dt} = \begin{cases} 0 & \text{In the absence of interactions with the universe (Galileo)} \\ \frac{1}{m} & \text{Multiplied by a functional form describing the interaction} \end{cases}$$

Some functions, fundamental ones

$$\mathbf{F} = \frac{-G m M}{r_{12}^2} \mathbf{e}_{12}$$

Others, phenomenological

$$F = -kx$$

## Newton's approach gave us more

We define the velocity.

$$\frac{dx}{dt} = v$$

$$\frac{dv}{dt} = \begin{cases} 0 & \text{In the absence of interactions with the universe (Galileo)} \\ \frac{1}{m} & \text{Multiplied by a functional form describing the interaction} \end{cases}$$

Some functions, fundamental ones

$$\mathbf{F} = \frac{-G m M}{r_{12}^2} \mathbf{e}_{12}$$

Others, phenomenological

$$F = -kx$$

To perform a prediction,  
we have to have a model relating  
the variables

$$\frac{dx}{dt} = y$$

$$\frac{dy}{dt} = f(x, y)$$

## More than a law, it is a program of how to do science

1. I measure a finite set of initial conditions.
2. I elucidate the law that prescribes the rate of variation of these variables (assuming that the law involves the values of the variables **at that moment**).
3. I integrate an ODE

It does NOT describe

1. Systems of equations with delays
2. Partial differential equations (although...)

$$\frac{dx(t)}{dt} = x(t) - \sigma x(t - \tau)^2$$

Even electromagnetism is excluded...

## Dynamical systems (in the framework of this course)

$$\frac{dx}{dt} = f(x) \quad x \in R^n$$

$$x(t = 0) = x_0$$

Under smoothness conditions for  $f(x)$  and  $df/dx$  in a neighborhood of  $x_0$ , the solution of the system exists and is unique. This means that each point (state) has a unique future (in geometrical terms, no auto-intersection of trajectories).

## Dynamical systems (in the framework of this course)


$$\frac{dx}{dt} = f(x) \quad x \in R^n$$

$$x(t = 0) = x_0$$




solving analytically these is  
usually impossible, therefore...


**Fixed points**



**Limit cycles**



**Chaotic attractors**



## Example: the relaxation oscillator

$$\frac{d^2x}{dt^2} + \mu + (x^2 - 1) \frac{dx}{dt} + kx = 0$$

$$\frac{d}{dt} \left( \frac{dx}{dt} + \frac{x^3}{3} - x \right) = -kx$$

$w$

$w$

$$\tilde{w} \equiv w/\mu$$




$$\frac{dx}{dt} = \mu (\tilde{w} - \frac{1}{3} x^3 + u)$$

$$\frac{d\tilde{w}}{dt} = -\frac{k}{\mu}(u)$$

## Example: the relaxation oscillator

$$\frac{d^2x}{dt^2} + \mu + (x^2 - 1) \frac{dx}{dt} + kx = 0$$

If  $\mu \gg 1$



$$\frac{dx}{dt} = \mu (\tilde{w} - \frac{1}{3} x^3 + u)$$
$$\frac{d\tilde{w}}{dt} = -\frac{k}{\mu}(u)$$

The “splash” affected  
all science, just as  
Laplace predicted


## A “Newtonian” approximation to neuroscience

$$C \frac{dV}{dt} = I - g_K n^4 (V - E_K) - g_{Na} m^3 h (V - E_{Na}) - a_+ (V - E_+) \quad \Big)$$

$$\frac{dn}{dt} = \alpha_n(V)(1 - n) - \beta_n(V)n$$


$$\frac{dm}{dt} = \alpha_m(V)(1 - m) - \beta_m(V)m$$

$$\frac{dh}{dt} = \alpha_h(V)(1 - h) - \beta_h(V)h$$



En que afecta al flujo el que el campo vector sea no lineal?


$$\frac{dx}{dt} = ax \left(1 - \frac{x}{N}\right).$$



La primera señal de que un sistema es no lineal es la coexistencia de puntos estacionarios aislados

$$\frac{dx}{dt} = r + x^2$$

$r \in \mathbb{R}$




La primera señal de que un sistema es no lineal es la coexistencia de puntos estacionarios aislados


$$\frac{dx}{dt} = r + x^2$$

$r \in \mathbb{R}$





Dependiendo del parámetro, existe un número diferente de puntos fijos. A tal cambio cualitativo se lo conoce como **bifurcación**. Notemos que un sistema no lineal puede tener un número mayor que uno de puntos fijos aislados.



A



B




## Nodo silla en ciclo límite




$$\frac{dr}{dt} = r(1 - r)$$

$$\frac{d\theta}{dt} = \omega - \cos\theta$$

Notemos que para la parte angular,



O sea, cuando  $\omega > 1$ , el sistema oscila. Las oscilaciones nacen con periodo infinito.



# Formas normales

$$\frac{dA_1}{dt} = A_2$$

$$\frac{dA_2}{dt} = (2\epsilon - A_1^2)A_2 - A_1$$

El primer termino de la segunda ecuación es una disipación (negativa si  $A_1$  es pequeña, y positiva si  $A_1$  es grande). El segundo término, una restitución.

Este sistema de ecuaciones, entonces, da lugar a una oscilación de relajación, como hemos visto en la segunda clase.

Miremos el problema cerca de donde ocurrirán las bifurcaciones, esto es, cerca de  $\epsilon \approx 0$ . Para simplificar, definamos

$$X = A_1 + iA_2$$

Lo que da lugar a la ecuación en variable compleja:

$$\frac{dX}{dt} = -iX - \frac{1}{8}(X^3 + X^2\bar{X} - X\bar{X}^2 - \bar{X}^3)$$

**Y la pregunta que nos vamos a hacer es si es posible encontrar un cambio de variables que elimine alguno de estos términos.**

Proponemos un cambio de variables implícito, con todos los términos cúbicos posibles, o sea:

$$X = Z + \alpha Z^3 + \beta Z^2 \bar{Z} + \gamma Z \bar{Z}^2 + \delta \bar{Z}^3$$

$$\begin{aligned} & (1 + 3\alpha Z^2 + 2\beta Z \bar{Z} + \gamma \bar{Z}^2) \frac{dZ}{dt} + (\beta Z^2 + 2\gamma Z \bar{Z} + 3\delta \bar{Z}^2) \frac{d\bar{Z}}{dt} = \\ & = -i(Z + \alpha Z^3 + \beta Z^2 \bar{Z} + \gamma Z \bar{Z}^2 + \delta \bar{Z}^3) - \frac{1}{8}(Z^3 + Z^2 \bar{Z} - Z \bar{Z}^2 - \bar{Z}^3) \end{aligned}$$

Y reemplazando  $\frac{d\bar{Z}}{dt}$  por  $i\bar{Z}$  (ya que va a entrar vía términos cúbicos, basta considerar la aproximación lineal), la ecuación queda:

$$\begin{aligned}\frac{dZ}{dt} = & -iZ + \left(2\alpha - \frac{1}{8}\right)Z^3 + \left(-\frac{1}{8}\right)Z^2\bar{Z} \\ & + \left(-2\gamma + \frac{1}{8}\right)Z\bar{Z}^2 + \left(-2\delta + \frac{1}{8}\right)\bar{Z}^3\end{aligned}$$

Notamos entonces, que podemos elegir  $\alpha, \gamma, \delta$  de modo que los términos con  $Z^3, Z\bar{Z}^2, \bar{Z}^3$  se vayan, quedando

$$\frac{dZ}{dt} = -iZ - \frac{1}{8}Z^2\bar{Z}$$

Esta es una expresión minimal. No hay cambio de coordenada posible que elimine a este termino.  
Notemos que es un termino tal que, si pensamos que linealmente

$$Z \sim e^{-it}$$

$$Z^2 \bar{Z} \sim e^{-2it} e^{it} = e^{-it}$$

Mientras que los términos eliminados van como

$$Z^3 \sim e^{-3it}, Z\bar{Z}^2 \sim e^{it}, \bar{Z}^3 \sim e^{3it}$$



Por eso decimos que el cambio no lineal de coordenadas dejo solo al termino **resonante**.

Esta estrategia ya la habíamos esbozado, cuando presentamos, para bifurcaciones en las primeras dos clases, ecuaciones sencillas, paradigmáticas, que representaran las propiedades cualitativas de un flujo, y sus cambios ante cambios de los parámetros. Por ejemplo:




Example

¿How do birds  
generate their  
songs?


# Example



# Example




# The program of physics (or interpretable science)




1. To identify the pertinent variables  
And how to relate them through a model

$$\left\{ \begin{array}{l} \frac{dx}{dt} = y, \\ \frac{dy}{dt} = (1/m) \left[ -k(x)x - b(y)y - cx^2y + a_{lab}p_s \left( \frac{\Delta a + 2\tau\tau}{a_{01} + x + \tau y} \right) \right]. \end{array} \right.$$

2. To study the solutions for different parameters




# The program of physics (or interpretable science)



Normal form

$$\left\{ \begin{array}{l} \frac{dx}{dt} = y \\ \frac{dy}{dt} = -a(t)g^2 - b(t)g^2x - g^2x^3 - gx^2y + g^2x^2 - gxy \end{array} \right.$$

1. To find the simplest models






## Normal form reduction




$$\left[ \begin{array}{l} \frac{dx}{dt} = y, \\ \frac{dy}{dt} = (1/m) \left[ -k(x)x - b(y)y - cx^2y + a_{\text{lab}} p_s \left( \frac{\Delta a + 2\tau\tau}{a_{01} + x + \tau y} \right) \right]. \end{array} \right] \xrightarrow{\text{Normal form reduction}} \left[ \begin{array}{l} \frac{dx}{dt} = y \\ \frac{dy}{dt} = -\alpha(t)\gamma^2 - \beta(t)\gamma^2 x - \gamma^2 x^3 - \gamma x^2 y + \gamma^2 x^2 - \gamma x y \end{array} \right]$$

Algorithmic procedure that allows you to reduce your nonlinear problem to a simpler one, close to a linear singularity. This simpler model will have less parameters, and **has been probably studied**

## Normal form reduction








**SNILC:** one of the ways oscillations arise in 2d systems

**How many?** The good news... not that many

STUDIES IN NONLINEARITY

# NONLINEAR DYNAMICS AND CHAOS



*With Applications to Physics, Biology,  
Chemistry, and Engineering*

STEVEN H. STROGATZ

John Guckenheimer  
Philip Holmes  
  
Applied  
Mathematical  
Sciences  
42  
  
Nonlinear  
Oscillations,  
Dynamical  
Systems,  
and Bifurcations  
of Vector Fields



## Newton's “mandate”

1. First, **understand the mechanisms**
2. Plug them into the machinery of **dynamical systems**
3. **Predict**

“Novelty” of the XX century:  
qualitative theory of dynamical systems and  
numerical integration as exploratory tool

But what if what we want to understand is extraordinarily complex?